

Continuity of Solution with Respect to Initial Data for One Class of Controlled Functional Differential Equation with Concentrated Delay Uniformly with Control Function

Mariam Khechikashvili

e-mail: mariam.khechikashvili@ens.tsu.edu.ge

Department of Mathematics, Iv. Javakishvili Tbilisi State University, 13 University St., 0186 Tbilisi, Georgia

For the differential equation

$$\dot{x}(t) = f(t, x(t), x(t - \tau), u(t))$$

with the initial condition

$$x(t) = \varphi_0(t) \quad , \quad t < t_{00}, \quad x(t_0) = x_{00}.$$

is proved theorem about continuous dependence of solution on initial data uniformly with respect to control functions $u(t) \in \Omega$, where Ω is the set of measurable control functions $u(t)$ with values in compact U . Under the initial data we mean the collection of the initial moment, delay parameter, initial function and initial vector. The theorem is proved by the scheme given in [1-3] .

References

- [1] T. Tadumadze , Some problems in the qualitative theory of optimal control.(Russian) *Tbilis. Gos. Univ.* , Tbilisi, 1983.
- [2] T. Tadumadze , Continuous dependence of solutions of delay functional differential equations on the right-hand side and initial data considering delay perturbations. *Georgin International Journal of Science and Technology*, **6**(4) (2014), 353-369.
- [3] T. Tadumadze, Variation formulas of solutions for functional differential equations with several constant delays and their applications in optimal control problems. *Mem . Differential Equations Math . Phys.* **70** (2017), 7-97.